

✖ WHO logo

[Home](#) [Countries](#) [Health topics](#) [Publications](#) [Research tools](#) [WHO sites](#) [FAQs](#)

WHO Statistical Information System (WHOSIS)

[WHO Statistics](#) -> [Selection screen](#) -> Display data

Core Health Indicators

the latest data from multiple WHO sources

These are the latest Core Health Indicators from WHO sources, including '*World Health Statistics 2006*' and '*The World Health Report, 2006 Edition*'.

[Make another selection](#) | [Group by indicators](#) | [Download selected data](#)

Algeria

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 69.0 |
| Life expectancy at birth (years) females (2) | | | | 72.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 59.7 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 61.6 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 153 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 124 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 41 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 39 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 35.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 20 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 140 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | <500 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 2 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 0 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 597.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 313.6 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 103.1 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 85.4 | | |
| Years of life lost to communicable diseases (%) (2) | | 49.7 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 30.1 | | |
| Years of life lost to injuries (%) (2) | | 20.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 48.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 11.9 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.9 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.5 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 13.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 5.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 20.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 2.5 | | | |
| Neonatal deaths due to infection (%) (2) | 25.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.5 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.4 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 9.6 | | | |
| Neonatal deaths due to preterm (%) (2) | 30.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.7 | | | |

Angola

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 38.0 |
| Life expectancy at birth (years) females (2) | | | | 42.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 31.6 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 35.1 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 591 |

| | | | | |
|--|-------|-------|--------|-------|
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 504 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 276 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 243 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 154.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 54 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,700 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 21,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 25 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 7 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 981.7 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 485.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 179.4 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 230.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 83.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.1 | | |
| Years of life lost to injuries (%) (2) | | 8.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 22.2 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 19.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.8 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 8.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 24.8 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 17.2 | | | |
| Neonatal deaths due to tetanus (%) (2) | 9.9 | | | |
| Neonatal deaths due to infection (%) (2) | 29.2 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 4.6 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.6 | | | |
| Neonatal deaths due to preterm (%) (2) | 21.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.7 | | | |

Benin

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|------|
| Life expectancy at birth (years) males (2) | | | | 52.0 |
| Life expectancy at birth (years) females (2) | | | | 53.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 43.4 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 44.5 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 388 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 350 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 152 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 153 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 90.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 38 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 850 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 5,800 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 14 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 2 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 852.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 432.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 153.7 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 116.2 | | |
| Years of life lost to communicable diseases (%) (2) | | 81.7 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 10.3 | | |
| Years of life lost to injuries (%) (2) | | 8.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 25.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 5.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 27.2 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.1 | | | |

| | | | |
|---|------|--|--|
| Deaths among children under five years of age due to injuries (%) (2) | 2.1 | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | |
| Neonatal deaths due to tetanus (%) (2) | 4.5 | | |
| Neonatal deaths due to infection (%) (2) | 33.4 | | |
| Neonatal deaths due to asphyxia (%) (2) | 20.0 | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.1 | | |
| Neonatal deaths due to congenital causes (%) (2) | 8.1 | | |
| Neonatal deaths due to preterm (%) (2) | 26.2 | | |
| Neonatal deaths due to other causes (%) (2) | 5.7 | | |

Botswana

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|------|
| Life expectancy at birth (years) males (2) | | | | 40.0 |
| Life expectancy at birth (years) females (2) | | | | 40.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 36.0 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 35.4 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 786 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 770 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 123 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 109 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 75.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 100 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 33,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 37 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 58 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 653.0 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 338.0 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 123.8 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 71.7 | | |
| Years of life lost to communicable diseases (%) (2) | | 93.1 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 4.3 | | |
| Years of life lost to injuries (%) (2) | | 2.6 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 40.3 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 53.8 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 1.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.3 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 2.9 | | | |
| Neonatal deaths due to infection (%) (2) | 22.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 20.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.4 | | | |
| Neonatal deaths due to preterm (%) (2) | 38.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 5.9 | | | |

Burkina Faso

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 47.0 |
| Life expectancy at birth (years) females (2) | | | | 48.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 34.9 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 36.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 472 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 410 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 193 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 191 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 97.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 36 | | | |

| | | | |
|--|-------|--------|----|
| Maternal mortality ratio (per 100 000 live births) (2) | 1,000 | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | 29,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | 38 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | 16 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 901.3 | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 458.7 | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 162.4 | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 148.9 | |
| Years of life lost to communicable diseases (%) (2) | | 86.8 | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.6 | |
| Years of life lost to injuries (%) (2) | | 6.6 | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 18.3 | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 4.0 | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.8 | | |
| Deaths among children under five years of age due to measles (%) (2) | 3.4 | | |
| Deaths among children under five years of age due to malaria (%) (2) | 20.3 | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.3 | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.5 | | |
| Deaths among children under five years of age due to other causes (%) (2) | 10.4 | | |
| Neonatal deaths due to tetanus (%) (2) | 11.8 | | |
| Neonatal deaths due to infection (%) (2) | 31.6 | | |
| Neonatal deaths due to asphyxia (%) (2) | 21.3 | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.4 | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.3 | | |
| Neonatal deaths due to preterm (%) (2) | 21.5 | | |
| Neonatal deaths due to other causes (%) (2) | 6.1 | | |

Burundi

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 42.0 |
| Life expectancy at birth (years) females (2) | | | | 47.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 33.4 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 36.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 593 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 457 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 196 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 184 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 114.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 41 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,000 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 25,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 59 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 33 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 843.4 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 438.7 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.4 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 300.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 80.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 7.2 | | |
| Years of life lost to injuries (%) (2) | | 11.9 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 23.3 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 8.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 8.4 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 22.8 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.8 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 14.6 | | | |
| Neonatal deaths due to tetanus (%) (2) | 9.9 | | | |
| Neonatal deaths due to infection (%) (2) | 31.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.9 | | | |

| | | | | |
|--|------|--|--|--|
| Neonatal deaths due to diarrhoea (%) (2) | 2.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.1 | | | |
| Neonatal deaths due to preterm (%) (2) | 19.7 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.8 | | | |

Cameroon

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|------|
| Life expectancy at birth (years) males (2) | | | | 50.0 |
| Life expectancy at birth (years) females (2) | | | | 51.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 41.1 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 444 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 432 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 156 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 143 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 87.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 730 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 49,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 19 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 12 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 848.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 435.7 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 149.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 118.4 | | |
| Years of life lost to communicable diseases (%) (2) | | 81.2 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.0 | | |
| Years of life lost to injuries (%) (2) | | 7.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 24.8 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 7.2 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.3 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 22.8 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.5 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 4.8 | | | |
| Neonatal deaths due to infection (%) (2) | 24.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.6 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.9 | | | |
| Neonatal deaths due to preterm (%) (2) | 27.7 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.2 | | | |

Cape Verde

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 67.0 |
| Life expectancy at birth (years) females (2) | | | | 71.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 58.8 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 62.9 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 209 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 139 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 38 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 35 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 27.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 10 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 150 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 35 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 1 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 691.8 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 356.2 | | |

| | | | | |
|--|------|-------|--|--|
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 127.3 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 38.9 | | |
| Years of life lost to communicable diseases (%) (2) | | 50.6 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 37.4 | | |
| Years of life lost to injuries (%) (2) | | 12.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 25.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 12.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.4 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 4.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 13.3 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.5 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 32.6 | | | |
| Neonatal deaths due to tetanus (%) (2) | 1.5 | | | |
| Neonatal deaths due to infection (%) (2) | 19.5 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 0.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 10.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 38.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.4 | | | |

Central African Republic

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 40.0 |
| Life expectancy at birth (years) females (2) | | | | 41.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 37.0 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 37.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 667 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 624 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 201 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 185 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 115.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 48 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,100 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 23,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 55 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 58 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 862.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 445.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 153.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 145.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 84.3 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.5 | | |
| Years of life lost to injuries (%) (2) | | 7.2 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 27.2 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 12.4 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 14.7 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.5 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 18.5 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 18.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 14.9 | | | |
| Neonatal deaths due to infection (%) (2) | 22.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.5 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.5 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.5 | | | |
| Neonatal deaths due to preterm (%) (2) | 23.4 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.4 | | | |

Chad

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 45.0 |
| Life expectancy at birth (years) females (2) | | | | 48.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 39.7 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 497 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 422 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 212 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 188 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 117.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 45 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,100 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 18,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 60 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 22 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 868.8 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 443.3 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 155.7 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 130.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 85.3 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.2 | | |
| Years of life lost to injuries (%) (2) | | 6.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 24.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 4.1 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 7.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 22.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 22.8 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.8 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 13.6 | | | |
| Neonatal deaths due to infection (%) (2) | 22.7 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.6 | | | |
| Neonatal deaths due to preterm (%) (2) | 25.3 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.8 | | | |

Comoros

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 62.0 |
| Life expectancy at birth (years) females (2) | | | | 67.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 53.9 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 55.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 254 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 182 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 76 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 64 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 52.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 29 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 480 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 7 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 0 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 736.5 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 381.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 128.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 83.2 | | |
| Years of life lost to communicable diseases (%) (2) | | 70.1 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 18.4 | | |
| Years of life lost to injuries (%) (2) | | 11.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 37.3 | | | |

| | | | | |
|--|------|--|--|--|
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 13.6 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 5.9 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 19.4 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 16.3 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.5 | | | |
| Neonatal deaths due to tetanus (%) (2) | 3.3 | | | |
| Neonatal deaths due to infection (%) (2) | 23.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.1 | | | |
| Neonatal deaths due to preterm (%) (2) | 34.8 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.8 | | | |

Congo

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|------|
| Life expectancy at birth (years) males (2) | | | | 53.0 |
| Life expectancy at birth (years) females (2) | | | | 55.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 45.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 47.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 442 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 390 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 113 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 103 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 79.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 32 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 510 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 9,700 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 52 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 18 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 761.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 392.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 134.1 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 146.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 78.8 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 10.7 | | |
| Years of life lost to injuries (%) (2) | | 10.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 30.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 9.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 11.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.6 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 25.7 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 13.6 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.6 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 3.8 | | | |
| Neonatal deaths due to infection (%) (2) | 17.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 27.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.2 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 8.3 | | | |
| Neonatal deaths due to preterm (%) (2) | 32.8 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.8 | | | |

Côte d'Ivoire

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 41.0 |
| Life expectancy at birth (years) females (2) | | | | 47.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 37.6 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 585 |

| | | | | |
|--|------|-------|--------|-------|
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 500 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 225 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 162 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 118.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 65 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 690 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 47,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 67 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 37 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 873.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 436.2 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 160.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 179.0 | | |
| Years of life lost to communicable diseases (%) (2) | | 78.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.1 | | |
| Years of life lost to injuries (%) (2) | | 10.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 34.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 5.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 14.8 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 20.5 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 19.6 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 7.7 | | | |
| Neonatal deaths due to infection (%) (2) | 32.3 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 20.2 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 4.6 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.6 | | | |
| Neonatal deaths due to preterm (%) (2) | 23.9 | | | |
| Neonatal deaths due to other causes (%) (2) | 5.7 | | | |

Democratic Republic of the Congo

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|---------|-------|
| Life expectancy at birth (years) males (2) | | | | 42.0 |
| Life expectancy at birth (years) females (2) | | | | 47.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 35.0 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 39.1 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 576 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 446 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 217 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 192 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 129.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 47 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 990 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 100,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 57 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 21 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 909.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 465.3 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 160.9 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 273.4 | | |
| Years of life lost to communicable diseases (%) (2) | | 81.7 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 7.4 | | |
| Years of life lost to injuries (%) (2) | | 10.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 25.7 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.7 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 16.9 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.1 | | | |

| | | | | |
|---|------|--|--|--|
| Deaths among children under five years of age due to injuries (%) (2) | 1.6 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 6.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 10.5 | | | |
| Neonatal deaths due to infection (%) (2) | 22.5 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.6 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 26.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.7 | | | |

Equatorial Guinea

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|-------|
| Life expectancy at birth (years) males (2) | | | | 42.0 |
| Life expectancy at birth (years) females (2) | | | | 44.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 44.7 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 46.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 577 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 522 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 213 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 195 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 123.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 880 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 30 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 27 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 863.7 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 438.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 155.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 144.1 | | |
| Years of life lost to communicable diseases (%) (2) | | 79.3 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.8 | | |
| Years of life lost to injuries (%) (2) | | 9.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 27.5 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 7.4 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 13.6 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 7.4 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 24.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 17.3 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 7.3 | | | |
| Neonatal deaths due to infection (%) (2) | 19.3 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 33.7 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.1 | | | |
| Neonatal deaths due to preterm (%) (2) | 20.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 9.6 | | | |

Eritrea

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 58.0 |
| Life expectancy at birth (years) females (2) | | | | 62.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 49.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 50.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 345 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 281 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 89 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 75 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 52.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 25 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 630 | | | |

| | | | | |
|--|------|-------|-------|----|
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 6,300 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 52 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 9 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 762.4 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 397.6 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 132.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 91.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 80.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.1 | | |
| Years of life lost to injuries (%) (2) | | 8.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 27.4 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 6.2 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 15.6 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 13.6 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 18.6 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 13.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 6.5 | | | |
| Neonatal deaths due to infection (%) (2) | 33.7 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.7 | | | |
| Neonatal deaths due to preterm (%) (2) | 20.4 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.3 | | | |

Ethiopia

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|---------|-------|
| Life expectancy at birth (years) males (2) | | | | 49.0 |
| Life expectancy at birth (years) females (2) | | | | 51.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 40.7 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 451 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 389 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 175 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 158 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 110.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 51 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 850 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 120,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 60 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 19 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 858.5 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 435.0 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.5 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 103.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 81.7 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 12.3 | | |
| Years of life lost to injuries (%) (2) | | 6.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 30.2 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.8 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.3 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.2 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 6.1 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 22.3 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.7 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 14.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 15.1 | | | |
| Neonatal deaths due to infection (%) (2) | 32.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.5 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 4.1 | | | |

| | | | | |
|--|------|--|--|--|
| Neonatal deaths due to congenital causes (%) (2) | 3.9 | | | |
| Neonatal deaths due to preterm (%) (2) | 15.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.4 | | | |

Gabon

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|------|
| Life expectancy at birth (years) males (2) | | | | 55.0 |
| Life expectancy at birth (years) females (2) | | | | 59.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 50.2 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 52.6 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 411 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 344 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 102 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 80 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 59.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 31 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 420 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 3,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 33 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 22 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 813.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 410.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 158.4 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 103.3 | | |
| Years of life lost to communicable diseases (%) (2) | | 72.3 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 18.3 | | |
| Years of life lost to injuries (%) (2) | | 9.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 35.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 10.1 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 8.8 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.4 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 28.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 10.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 5.3 | | | |
| Neonatal deaths due to infection (%) (2) | 23.6 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 9.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 30.7 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.4 | | | |

Gambia

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 55.0 |
| Life expectancy at birth (years) females (2) | | | | 59.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 48.5 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 50.5 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 344 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 263 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 129 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 115 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 89.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 46 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 540 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 600 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 38 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 2 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 805.0 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 412.6 | | |

| | | | |
|--|-------|--|--|
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | 144.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | 108.8 | | |
| Years of life lost to communicable diseases (%) (2) | 75.0 | | |
| Years of life lost to non-communicable diseases (%) (2) | 15.2 | | |
| Years of life lost to injuries (%) (2) | 9.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 36.6 | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 1.3 | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 12.2 | | |
| Deaths among children under five years of age due to measles (%) (2) | 2.5 | | |
| Deaths among children under five years of age due to malaria (%) (2) | 29.4 | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 15.5 | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.6 | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | |
| Neonatal deaths due to tetanus (%) (2) | 5.0 | | |
| Neonatal deaths due to infection (%) (2) | 37.2 | | |
| Neonatal deaths due to asphyxia (%) (2) | 21.1 | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.0 | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.5 | | |
| Neonatal deaths due to preterm (%) (2) | 22.2 | | |
| Neonatal deaths due to other causes (%) (2) | 6.0 | | |

Ghana

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|------|
| Life expectancy at birth (years) males (2) | | | | 56.0 |
| Life expectancy at birth (years) females (2) | | | | 58.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 49.2 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 50.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 349 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 319 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 113 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 111 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 68.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 27 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 540 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 30,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 40 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 10 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 785.8 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 404.2 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 137.7 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 96.7 | | |
| Years of life lost to communicable diseases (%) (2) | | 74.4 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 15.6 | | |
| Years of life lost to injuries (%) (2) | | 10.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 28.5 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 5.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 12.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 2.9 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 33.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 14.6 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 2.5 | | | |
| Neonatal deaths due to infection (%) (2) | 29.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 26.5 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.5 | | | |
| Neonatal deaths due to preterm (%) (2) | 24.7 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.5 | | | |

Guinea

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|-------|
| Life expectancy at birth (years) males (2) | | | | 52.0 |
| Life expectancy at birth (years) females (2) | | | | 55.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 43.9 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 45.6 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 364 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 319 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 160 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 150 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 101.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 48 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 740 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 9,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 44 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 12 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 853.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 432.2 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 155.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 147.4 | | |
| Years of life lost to communicable diseases (%) (2) | | 80.0 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 10.7 | | |
| Years of life lost to injuries (%) (2) | | 9.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 28.8 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 2.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 5.5 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 24.5 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 20.9 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 10.9 | | | |
| Neonatal deaths due to infection (%) (2) | 31.2 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.0 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.4 | | | |
| Neonatal deaths due to preterm (%) (2) | 19.9 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.5 | | | |

Guinea-Bissau

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|------|-------|
| Life expectancy at birth (years) males (2) | | | | 45.0 |
| Life expectancy at birth (years) females (2) | | | | 48.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 39.6 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.5 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 482 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 413 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 212 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 194 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 126.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 48 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,100 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 33 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 9 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 882.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 449.2 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 159.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 137.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 85.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.0 | | |
| Years of life lost to injuries (%) (2) | | 6.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 24.1 | | | |

| | | | | |
|--|------|--|--|--|
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 2.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.6 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 3.4 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 21.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.4 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 5.5 | | | |
| Neonatal deaths due to tetanus (%) (2) | 12.9 | | | |
| Neonatal deaths due to infection (%) (2) | 31.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 20.7 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.3 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.0 | | | |
| Neonatal deaths due to preterm (%) (2) | 20.1 | | | |
| Neonatal deaths due to other causes (%) (2) | 5.9 | | | |

Kenya

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|---------|------|
| Life expectancy at birth (years) males (2) | | | | 51.0 |
| Life expectancy at birth (years) females (2) | | | | 50.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 44.1 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 44.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 477 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 502 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 129 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 110 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 78.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 29 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,000 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 150,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 90 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 43 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 782.5 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 401.5 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 139.3 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 95.3 | | |
| Years of life lost to communicable diseases (%) (2) | | 81.4 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.1 | | |
| Years of life lost to injuries (%) (2) | | 7.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 24.2 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 14.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 3.2 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 13.6 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 19.9 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.7 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 5.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 2.7 | | | |
| Neonatal deaths due to infection (%) (2) | 25.7 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 27.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 26.0 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.9 | | | |

Lesotho

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 39.0 |
| Life expectancy at birth (years) females (2) | | | | 44.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 29.6 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 33.2 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 845 |

| | | | | |
|--|------|-------|--------|------|
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 728 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 87 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 76 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 55.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 28 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 550 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 29,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 41 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 58 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 784.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 404.3 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 139.3 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 87.6 | | |
| Years of life lost to communicable diseases (%) (2) | | 90.1 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.6 | | |
| Years of life lost to injuries (%) (2) | | 3.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 32.8 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 56.2 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 3.9 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 4.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 4.6 | | | |
| Neonatal deaths due to infection (%) (2) | 22.5 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.5 | | | |
| Neonatal deaths due to preterm (%) (2) | 30.0 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.2 | | | |

Liberia

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|-------|
| Life expectancy at birth (years) males (2) | | | | 39.0 |
| Life expectancy at birth (years) females (2) | | | | 44.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 33.6 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 37.0 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 596 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 477 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 249 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 220 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 157.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 66 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 760 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 7,200 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 46 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 27 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 955.3 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 485.3 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 169.4 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 269.6 | | |
| Years of life lost to communicable diseases (%) (2) | | 83.4 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.9 | | |
| Years of life lost to injuries (%) (2) | | 9.7 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 29.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.3 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 18.9 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.0 | | | |

| | | | | |
|---|------|--|--|--|
| Deaths among children under five years of age due to injuries (%) (2) | 1.7 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 18.6 | | | |
| Neonatal deaths due to infection (%) (2) | 27.0 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 16.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 5.0 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 3.3 | | | |
| Neonatal deaths due to preterm (%) (2) | 25.0 | | | |
| Neonatal deaths due to other causes (%) (2) | 4.7 | | | |

Madagascar

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|------|
| Life expectancy at birth (years) males (2) | | | | 55.0 |
| Life expectancy at birth (years) females (2) | | | | 59.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 47.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 49.9 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 338 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 270 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 128 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 117 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 76.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 33 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 550 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 7,500 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 37 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 6 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 836.7 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 429.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 147.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 112.0 | | |
| Years of life lost to communicable diseases (%) (2) | | 78.6 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 12.4 | | |
| Years of life lost to injuries (%) (2) | | 8.9 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 25.6 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 1.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.9 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 5.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 20.1 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 20.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 8.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 6.2 | | | |
| Neonatal deaths due to infection (%) (2) | 26.3 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 24.7 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.2 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.2 | | | |
| Neonatal deaths due to preterm (%) (2) | 27.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.0 | | | |

Malawi

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|-------|
| Life expectancy at birth (years) males (2) | | | | 41.0 |
| Life expectancy at birth (years) females (2) | | | | 41.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 35.0 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 34.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 663 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 638 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 179 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 172 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 109.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |

| | | | |
|--|-------|--------|----|
| Maternal mortality ratio (per 100 000 live births) (2) | 1,800 | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | 84,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | 48 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | 49 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 835.0 | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 429.6 | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 150.5 | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 105.3 | |
| Years of life lost to communicable diseases (%) (2) | | 89.3 | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.2 | |
| Years of life lost to injuries (%) (2) | | 4.5 | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 21.7 | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 14.0 | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.1 | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.3 | | |
| Deaths among children under five years of age due to malaria (%) (2) | 14.1 | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 22.6 | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.7 | | |
| Deaths among children under five years of age due to other causes (%) (2) | 7.6 | | |
| Neonatal deaths due to tetanus (%) (2) | 5.3 | | |
| Neonatal deaths due to infection (%) (2) | 28.7 | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.8 | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.8 | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.0 | | |
| Neonatal deaths due to preterm (%) (2) | 26.9 | | |
| Neonatal deaths due to other causes (%) (2) | 6.5 | | |

Mali

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 44.0 |
| Life expectancy at birth (years) females (2) | | | | 47.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 37.5 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 38.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 490 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 414 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 230 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 208 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 121.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 55 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,200 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 12,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 62 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 11 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 909.0 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 456.0 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 165.9 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 145.4 | | |
| Years of life lost to communicable diseases (%) (2) | | 85.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 7.7 | | |
| Years of life lost to injuries (%) (2) | | 6.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 25.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 1.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.3 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 16.9 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.9 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 5.9 | | | |
| Neonatal deaths due to tetanus (%) (2) | 11.4 | | | |
| Neonatal deaths due to infection (%) (2) | 31.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.1 | | | |

| | | | | |
|--|------|--|--|--|
| Neonatal deaths due to diarrhoea (%) (2) | 4.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.7 | | | |
| Neonatal deaths due to preterm (%) (2) | 19.3 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.3 | | | |

Mauritania

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 55.0 |
| Life expectancy at birth (years) females (2) | | | | 60.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 42.8 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 46.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 325 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 246 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 134 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 115 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 78.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 70 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,000 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | <500 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 57 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 3 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 884.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 451.1 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 157.5 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 137.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 79.2 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.7 | | |
| Years of life lost to injuries (%) (2) | | 9.1 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 39.4 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 0.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 1.7 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 12.2 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 22.3 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.9 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 5.9 | | | |
| Neonatal deaths due to tetanus (%) (2) | 21.4 | | | |
| Neonatal deaths due to infection (%) (2) | 29.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 17.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 5.5 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 16.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 4.9 | | | |

Mauritius

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 69.0 |
| Life expectancy at birth (years) females (2) | | | | 75.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 60.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 64.6 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 217 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 112 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 17 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 14 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 12.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 12 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 24 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 11 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 0 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 700.9 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 433.9 | | |

| | | | | |
|--|------|------|--|--|
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 78.9 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 41.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 11.4 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 75.5 | | |
| Years of life lost to injuries (%) (2) | | 13.1 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 66.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 1.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 3.9 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 5.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 23.6 | | | |
| Neonatal deaths due to tetanus (%) (2) | 0.0 | | | |
| Neonatal deaths due to infection (%) (2) | 8.4 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 15.2 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 0.4 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 13.1 | | | |
| Neonatal deaths due to preterm (%) (2) | 59.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 3.5 | | | |

Mozambique

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|---------|-------|
| Life expectancy at birth (years) males (2) | | | | 44.0 |
| Life expectancy at birth (years) females (2) | | | | 46.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 36.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 37.5 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 627 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 549 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 154 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 150 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 102.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 48 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,000 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 110,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 62 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 67 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 720.0 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 370.6 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 124.5 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 66.0 | | |
| Years of life lost to communicable diseases (%) (2) | | 91.1 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.6 | | |
| Years of life lost to injuries (%) (2) | | 2.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 29.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 12.9 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 18.9 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.2 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 10.1 | | | |
| Neonatal deaths due to infection (%) (2) | 37.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 18.8 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 5.0 | | | |
| Neonatal deaths due to preterm (%) (2) | 19.1 | | | |
| Neonatal deaths due to other causes (%) (2) | 5.4 | | | |

Namibia

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|------|
| Life expectancy at birth (years) males (2) | | | | 52.0 |
| Life expectancy at birth (years) females (2) | | | | 55.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 42.9 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 43.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 548 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 489 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 70 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 57 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 42.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 25 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 300 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 16,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 44 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 41 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 754.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 385.4 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 93.2 | | |
| Years of life lost to communicable diseases (%) (2) | | 83.3 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 10.4 | | |
| Years of life lost to injuries (%) (2) | | 6.3 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 38.5 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 53.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 2.2 | | | |
| Neonatal deaths due to infection (%) (2) | 18.9 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 24.0 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 8.7 | | | |
| Neonatal deaths due to preterm (%) (2) | 37.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.8 | | | |

Niger

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|-------|-------|
| Life expectancy at birth (years) males (2) | | | | 42.0 |
| Life expectancy at birth (years) females (2) | | | | 41.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 35.8 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 35.2 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 506 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 478 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 256 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 262 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 152.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 43 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,600 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 4,800 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 31 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 3 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 916.5 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 456.1 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 169.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 162.6 | | |
| Years of life lost to communicable diseases (%) (2) | | 87.0 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.7 | | |
| Years of life lost to injuries (%) (2) | | 6.3 | | |

| | | | | |
|--|------|--|--|--|
| Deaths among children under five years of age due to neonatal causes (%) (2) | 16.7 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 0.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 19.8 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 7.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 14.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 25.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.4 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 14.8 | | | |
| Neonatal deaths due to tetanus (%) (2) | 16.0 | | | |
| Neonatal deaths due to infection (%) (2) | 30.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.2 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.7 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 4.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 16.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.6 | | | |

Nigeria

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|---------|-------|
| Life expectancy at birth (years) males (2) | | | | 45.0 |
| Life expectancy at birth (years) females (2) | | | | 46.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 41.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 41.8 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 513 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 478 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 198 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 195 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 103.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 53 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 800 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 310,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 56 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 27 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 889.3 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 451.6 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 156.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 131.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 82.6 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 10.1 | | |
| Years of life lost to injuries (%) (2) | | 7.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 26.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 5.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 15.7 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 24.1 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 20.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.9 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.8 | | | |
| Neonatal deaths due to tetanus (%) (2) | 10.3 | | | |
| Neonatal deaths due to infection (%) (2) | 23.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.9 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.5 | | | |
| Neonatal deaths due to preterm (%) (2) | 23.4 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.3 | | | |

Rwanda

| Indicator | 2000 | 2002 | 2003 | 2004 |
|---|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 44.0 |
| Life expectancy at birth (years) females (2) | | | | 47.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 36.4 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 40.2 | | |

| | | | | |
|--|-------|-------|--------|-------|
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 518 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 435 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 211 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 195 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 118.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 45 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,400 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 22,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 69 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 33 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 831.4 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 424.9 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 150.2 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 125.7 | | |
| Years of life lost to communicable diseases (%) (2) | | 84.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.2 | | |
| Years of life lost to injuries (%) (2) | | 6.9 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 21.7 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 5.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 18.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 1.6 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 4.6 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 23.2 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.8 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 23.7 | | | |
| Neonatal deaths due to tetanus (%) (2) | 3.4 | | | |
| Neonatal deaths due to infection (%) (2) | 29.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 27.3 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.2 | | | |
| Neonatal deaths due to preterm (%) (2) | 23.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.8 | | | |

Sao Tome and Principe

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|------|------|
| Life expectancy at birth (years) males (2) | | | | 57.0 |
| Life expectancy at birth (years) females (2) | | | | 60.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 54.2 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 54.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 301 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 236 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 122 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 114 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 75.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 38 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 28 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 0 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 764.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 395.5 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 133.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 86.9 | | |
| Years of life lost to communicable diseases (%) (2) | | 67.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 20.8 | | |
| Years of life lost to injuries (%) (2) | | 11.7 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 32.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 3.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.0 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 4.8 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.6 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.2 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.5 | | | |

| | | | | |
|---|------|--|--|--|
| Deaths among children under five years of age due to other causes (%) (2) | 18.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 1.6 | | | |
| Neonatal deaths due to infection (%) (2) | 19.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 24.0 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.0 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 11.8 | | | |
| Neonatal deaths due to preterm (%) (2) | 34.7 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.8 | | | |

Senegal

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|-------|------|
| Life expectancy at birth (years) males (2) | | | | 54.0 |
| Life expectancy at birth (years) females (2) | | | | 57.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 47.1 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 48.9 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 358 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 288 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 141 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 132 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 78.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 31 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 690 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 3,500 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 49 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 3 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 831.6 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 426.3 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.3 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 125.2 | | |
| Years of life lost to communicable diseases (%) (2) | | 75.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 13.2 | | |
| Years of life lost to injuries (%) (2) | | 10.9 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 22.8 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 1.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 8.1 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 27.6 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 20.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.6 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.2 | | | |
| Neonatal deaths due to tetanus (%) (2) | 6.1 | | | |
| Neonatal deaths due to infection (%) (2) | 32.2 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.7 | | | |
| Neonatal deaths due to preterm (%) (2) | 24.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.4 | | | |

Seychelles

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 67.0 |
| Life expectancy at birth (years) females (2) | | | | 78.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 57.4 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 64.9 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 232 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 83 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 14 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 13 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 12.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 9 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 6 |

| | | | | |
|--|------|-------|--|---|
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 0 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 657.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 335.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 130.7 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 68.6 | | |
| Years of life lost to communicable diseases (%) (2) | | 15.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 63.7 | | |
| Years of life lost to injuries (%) (2) | | 20.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 27.2 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 10.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 12.3 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 50.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 0.8 | | | |
| Neonatal deaths due to infection (%) (2) | 16.2 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.1 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 0.8 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 11.4 | | | |
| Neonatal deaths due to preterm (%) (2) | 42.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.3 | | | |

Sierra Leone

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|---------|------|-------|
| Life expectancy at birth (years) males (2) | | | | 37.0 |
| Life expectancy at birth (years) females (2) | | | | 40.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 27.2 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 29.9 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 579 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 497 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 296 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 269 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 165.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 56 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 2,000 | | | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 92 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 13 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 1,016.7 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 514.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 181.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 250.1 | | |
| Years of life lost to communicable diseases (%) (2) | | 85.7 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.5 | | |
| Years of life lost to injuries (%) (2) | | 7.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 21.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 1.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 19.7 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 5.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 12.4 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 25.5 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 12.7 | | | |
| Neonatal deaths due to tetanus (%) (2) | 21.2 | | | |
| Neonatal deaths due to infection (%) (2) | 28.3 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 16.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 4.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 3.4 | | | |
| Neonatal deaths due to preterm (%) (2) | 21.3 | | | |
| Neonatal deaths due to other causes (%) (2) | 4.8 | | | |

South Africa

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|---------|------|
| Life expectancy at birth (years) males (2) | | | | 47.0 |
| Life expectancy at birth (years) females (2) | | | | 49.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 43.3 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 45.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 667 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 598 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 72 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 62 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 54.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 21 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 230 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 370,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 56 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 78 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 808.3 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 409.9 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 154.1 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 119.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 77.0 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 14.6 | | |
| Years of life lost to injuries (%) (2) | | 8.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 35.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 57.1 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 0.8 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.0 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 0.9 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 5.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 1.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 1.4 | | | |
| Neonatal deaths due to infection (%) (2) | 20.8 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 23.2 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 1.4 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 10.4 | | | |
| Neonatal deaths due to preterm (%) (2) | 36.3 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.6 | | | |

Swaziland

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 36.0 |
| Life expectancy at birth (years) females (2) | | | | 39.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 33.2 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 35.2 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 823 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 741 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 163 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 150 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 102.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 38 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 370 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 17,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 93 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 176 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 731.5 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 363.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 161.9 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 71.8 | | |
| Years of life lost to communicable diseases (%) (2) | | 91.2 | | |

| | | | | |
|--|------|-----|--|--|
| Years of life lost to non-communicable diseases (%) (2) | | 5.3 | | |
| Years of life lost to injuries (%) (2) | | 3.5 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 26.8 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 47.0 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 9.6 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 0.2 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.2 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 11.8 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 3.8 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.5 | | | |
| Neonatal deaths due to tetanus (%) (2) | 1.8 | | | |
| Neonatal deaths due to infection (%) (2) | 25.3 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 26.1 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.6 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 8.3 | | | |
| Neonatal deaths due to preterm (%) (2) | 28.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.4 | | | |

Togo

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|------|
| Life expectancy at birth (years) males (2) | | | | 52.0 |
| Life expectancy at birth (years) females (2) | | | | 56.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 43.5 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 45.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 401 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 327 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 151 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 128 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 79.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 570 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 10,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 77 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 21 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 831.2 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 426.5 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.6 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 116.7 | | |
| Years of life lost to communicable diseases (%) (2) | | 79.4 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 11.8 | | |
| Years of life lost to injuries (%) (2) | | 8.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 29.0 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 5.8 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 13.8 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 6.6 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 25.3 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 17.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.5 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 6.5 | | | |
| Neonatal deaths due to infection (%) (2) | 29.5 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 22.9 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.5 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.7 | | | |
| Neonatal deaths due to preterm (%) (2) | 25.3 | | | |
| Neonatal deaths due to other causes (%) (2) | 6.5 | | | |

Uganda

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 48.0 |
| Life expectancy at birth (years) females (2) | | | | 51.0 |

| | | | | |
|--|------|-------|--------|------|
| Healthy life expectancy (HALE) at birth (years) males (2) | | 41.7 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 43.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 525 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 446 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 144 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 132 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 81.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 32 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 880 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 78,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 71 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 22 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 824.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 422.1 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 146.4 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 153.5 | | |
| Years of life lost to communicable diseases (%) (2) | | 83.9 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 7.9 | | |
| Years of life lost to injuries (%) (2) | | 8.2 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 23.6 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 7.7 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.2 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 3.0 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 23.1 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 2.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 4.9 | | | |
| Neonatal deaths due to infection (%) (2) | 26.2 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 26.7 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.2 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.9 | | | |
| Neonatal deaths due to preterm (%) (2) | 25.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.6 | | | |

United Republic of Tanzania

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|-------|-------|---------|------|
| Life expectancy at birth (years) males (2) | | | | 47.0 |
| Life expectancy at birth (years) females (2) | | | | 49.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 40.0 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 40.7 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 551 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 524 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 134 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 117 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 78.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 43 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,500 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 160,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 47 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 31 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 847.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 434.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 150.8 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 114.6 | | |
| Years of life lost to communicable diseases (%) (2) | | 85.5 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 8.2 | | |
| Years of life lost to injuries (%) (2) | | 6.4 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 26.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 9.3 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 16.8 | | | |

| | | | | |
|---|------|--|--|--|
| Deaths among children under five years of age due to measles (%) (2) | 1.3 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 22.7 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.1 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 2.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.0 | | | |
| Neonatal deaths due to tetanus (%) (2) | 3.4 | | | |
| Neonatal deaths due to infection (%) (2) | 28.7 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 26.6 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.1 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 7.2 | | | |
| Neonatal deaths due to preterm (%) (2) | 23.5 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.6 | | | |

Zambia

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|-------|--------|-------|
| Life expectancy at birth (years) males (2) | | | | 40.0 |
| Life expectancy at birth (years) females (2) | | | | 40.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 34.8 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 35.0 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 683 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 656 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 190 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 173 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 104.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 40 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 750 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 89,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 70 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 68 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 700.1 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 358.7 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 122.0 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 58.0 | | |
| Years of life lost to communicable diseases (%) (2) | | 92.0 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.0 | | |
| Years of life lost to injuries (%) (2) | | 2.0 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 22.9 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 16.1 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 17.5 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 1.2 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 19.4 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 21.8 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.0 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.1 | | | |
| Neonatal deaths due to tetanus (%) (2) | 3.8 | | | |
| Neonatal deaths due to infection (%) (2) | 28.1 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.4 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 3.0 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 6.9 | | | |
| Neonatal deaths due to preterm (%) (2) | 25.6 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.2 | | | |

Zimbabwe

| Indicator | 2000 | 2002 | 2003 | 2004 |
|--|------|------|------|------|
| Life expectancy at birth (years) males (2) | | | | 37.0 |
| Life expectancy at birth (years) females (2) | | | | 34.0 |
| Healthy life expectancy (HALE) at birth (years) males (2) | | 33.8 | | |
| Healthy life expectancy (HALE) at birth (years) females (2) | | 33.3 | | |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) males (2) | | | | 857 |
| Probability of dying (per 1 000 population) between 15 and 60 years (adult mortality rate) females (2) | | | | 849 |
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) males (2) | | | | 136 |

| | | | | |
|--|-------|-------|---------|------|
| Probability of dying (per 1 000 population) under five years of age (under-5 mortality rate) females (2) | | | | 121 |
| Infant mortality rate (per 1 000 live births) (2) | | | | 78.0 |
| Neonatal mortality rate (per 1 000 live births) (2) | 33 | | | |
| Maternal mortality ratio (per 100 000 live births) (2) | 1,100 | | | |
| Deaths due to HIV/AIDS (per 100 000 population per year) (2) | | | 170,000 | |
| Deaths due to tuberculosis among HIV-negative people (per 100 000 population) (2) | | | | 62 |
| Deaths due to tuberculosis among HIV-positive people (per 100 000 population) (2) | | | | 89 |
| Age-standardized mortality rate for non-communicable diseases (per 100 000 population) (2) | | 685.4 | | |
| Age-standardized mortality rate for cardiovascular diseases (per 100 000 population) (2) | | 346.8 | | |
| Age-standardized mortality rate for cancer (per 100 000 population) (2) | | 121.8 | | |
| Age-standardized mortality rate for injuries (per 100 000 population) (2) | | 103.3 | | |
| Years of life lost to communicable diseases (%) (2) | | 89.6 | | |
| Years of life lost to non-communicable diseases (%) (2) | | 6.6 | | |
| Years of life lost to injuries (%) (2) | | 3.8 | | |
| Deaths among children under five years of age due to neonatal causes (%) (2) | 28.1 | | | |
| Deaths among children under five years of age due to HIV/AIDS (%) (2) | 40.6 | | | |
| Deaths among children under five years of age due to diarrhoeal diseases (%) (2) | 12.1 | | | |
| Deaths among children under five years of age due to measles (%) (2) | 2.9 | | | |
| Deaths among children under five years of age due to malaria (%) (2) | 0.2 | | | |
| Deaths among children under five years of age due to pneumonia (%) (2) | 14.7 | | | |
| Deaths among children under five years of age due to injuries (%) (2) | 1.2 | | | |
| Deaths among children under five years of age due to other causes (%) (2) | 0.3 | | | |
| Neonatal deaths due to tetanus (%) (2) | 1.9 | | | |
| Neonatal deaths due to infection (%) (2) | 19.6 | | | |
| Neonatal deaths due to asphyxia (%) (2) | 25.8 | | | |
| Neonatal deaths due to diarrhoea (%) (2) | 2.0 | | | |
| Neonatal deaths due to congenital causes (%) (2) | 9.2 | | | |
| Neonatal deaths due to preterm (%) (2) | 34.2 | | | |
| Neonatal deaths due to other causes (%) (2) | 7.3 | | | |